

Certificate of Analysis

Custom Services for

Human induced Pluripotent Stem Cell (iPSC) Line: **OR00007*B**

Diagnosis	Developmental and Epileptic Encephalopathy 2
Parental cell line mutation	CDKL5; c.1412delA (p. Asp471fs)
Parental cell type, cell line ID	Fibroblast,
Sex	Female
Reprogramming method	Sendai viral vectors containing OCT4, SOX2, KLF4, and CMYC
Passage number at freeze	P8
Culture media	mTeSR1™
Feeder or Matrix substrate	Matrigel®
Recommended passage method and split ratio	Versene; 1:8 every 5-7 days*
iPSC line establishment publication(s)	

The following testing specifications have been met for this product lot:

Test Description	Test Method	Test Specification	Result
Post-Thaw Cell Viability	Colony doubling	Colony formation and diameter doubling within 5 days	Pass
Sterility	Growth on agar and broth	Negative	Pass
Mycoplasma	qRT-PCR	Negative	Pass
Alkaline Phosphatase Staining	Cell staining	> 80% cells with positive staining	Pass
Identity Match	STR (THO-1, D22S417, D10S526, vWA31, D5S592, and FES/FPS)	Match cell line	Pass
Genomic Integration of Episomal Plasmid	Genomic PCR using plasmid specific primers and endogenous FBXO1 control	No plasmid specific sequence amplified using 100ng gDNA template	N/A
Detection of Sendai Virus Genome and Transgene	qRT-PCR using SEV specific primers	No detection of SEV genome or transgenes	N/A
Surface Antigen Expression of Stem Cell Markers	Immunostaining and flow cytometric detection	> 80% expression of SSEA4	Pass
Differentiation Potential	Embryoid Body (EB) formation and gene expression	Minimal of 1 gene per germ layer expressed 2 fold or higher	Pass
Cytogenomics	G-banding, Affymetrix Human SNP Array 6.0	46,XX[20].arr(1-22,X)x2	Pass

Note: *Recover into 2 wells of a 6 well plate.

Christine Grandizio 08/11/2020

 Technician, Stem Cell Laboratory Date

Christine Grandizio 08/14/2020

 Manager, Stem Cell Laboratory Date

Disclaimer: iPSC lines distributed by Coriell Institute for Medical Research may differ from those in the submitter's laboratory.

Form 1701-10 Rev D-110519: Custom Services Certificate of Analysis OR00007*B

Post-Thaw Cell Viability

One distribution lot vial of the cell line was thawed and placed in culture. Cultures were observed daily. Colonies were photographed upon first appearance, then 3 days later. Colonies must double in diameter within 5 days. The area for 5 colonies was measured using CellSens software on the Olympus IX50 microscope at 40x magnification. The average area is reported here.

Day	Average area (μm^2)
1	6,617
4	212,276

Colony area increased by 32 fold.





Figure 1A. Colonies post thaw (Day 1)

Figure 1B. Colonies 3 days after first observation (Day 4)

Alkaline Phosphatase Staining

Cells were stained using the StemTAG™ Alkaline Phosphatase Staining Kit from CellBiolabs, Inc.




Figure 2. iPSC colonies showing alkaline phosphatase activity



Surface Antigen Expression of Stem Cell Markers

Undifferentiated cells are stained for stage specific embryonic antigen 4 (SSEA4) which is expressed on the surface of undifferentiated human pluripotent stem cells. Cells were analyzed using the MACSQuant Flow Cytometer by Miltenyi Biotec. More than 80% of cells should stain with antibodies specific for SSEA4.




Figure 3. Representative histogram of SSEA4 positive population. Histogram is an overlay of isotype stained control (red) and SSEA4 positive population (blue).



Surface Antigen Expression of Stem Cell Markers

Undifferentiated cells are stained for octamer-binding transcription factor 4 (OCT4) which is involved in self-renewal of undifferentiated human pluripotent stem cells to maintain pluripotency. Cells were analyzed using the MACSQuant Flow Cytometer by Miltenyi Biotec. More than 80% of cells should stain with antibodies specific for OCT4.





Figure 4. Representative histogram of OCT4 positive population. Histogram is an overlay of isotype stained control (red) and OCT4 positive population (blue).

Differentiation Potential

Cells are differentiated by embryoid body (EB) formation to assess pluripotency. RNA is extracted and gene expression is measured by quantitative RT-PCR. Ct values are adjusted to the endogenous housekeeping gene GAPDH. Relative gene expression is shown as the fold difference in expression compared to undifferentiated cells. Expression of at least one gene per germ layer should increase by 2 fold or higher.



Gene	Fold change	Gene	Fold change	Gene	Fold change	Gene	Fold change
OCT4	0	PAX6	27	T	70	AFP	766614
SOX2	0	NES	0	RUNX1	46	SOX17	409
NANOG	0	TP63	177	DES	2	FOXA2	244
GDF3	0	KRT14	79	PECAM1	212	SOX7	27
REXO1	0	NOG	193	TAL1	21		

Figure 4. Fold change in expression of pluripotency genes and tri-lineage specific genes

Note: Negative values are set as 0. Calculations are performed using the $2^{-\Delta\Delta CT}$ method. (Livak KJ, Schmittgen TD. *Methods*. 2001 Dec;25(4):402-8.PMID:11846609)

Cytogenomics

Microarray	Affymetrix Human SNP Array 6.0
Cytogenetic Banding Technique	G-banding
Passage at Analysis	P10
Metaphase Cells Counted	20
Metaphase Cells Analyzed	5
Metaphase Cells Karyotyped	5
Short ISCN	46,XX[20].arr(1-22,X)x2




Figure 5. G-banding karyogram